


Atomic orbitals of finite range as basis sets



Javier Junquera



UNIVERSIDAD DE CANTABRIA

Most important reference followed in this lecture

phys. stat. sol. (b) **215**, 809 (1999)

Subject classification: 71.15.Mb; 71.15.Fv; 71.24.+q; S1.3; S5; S5.11

Linear-Scaling ab-initio Calculations for Large and Complex Systems

E. ARTACHO¹) (a), D. SÁNCHEZ-PORTAL (b), P. ORDEJÓN (c), A. GARCÍA (d),
and J. M. SOLER (e)

PHYSICAL REVIEW B, VOLUME 64, 235111

Numerical atomic orbitals for linear-scaling calculations

Javier Junquera,¹ Óscar Paz,¹ Daniel Sánchez-Portal,^{2,3} and Emilio Artacho⁴

PHYSICAL REVIEW B **66**, 205101 (2002)

Systematic generation of finite-range atomic basis sets for linear-scaling calculations

Eduardo Anglada,^{1,2} José M. Soler,¹ Javier Junquera,³ and Emilio Artacho⁴

...in previous chapters:

the many body problem reduced to a problem of independent particles

One particle Kohn-Sham equation

$$\left[-\frac{1}{2} \nabla^2 + V_{eff}^{\sigma}(\vec{r}) \right] \psi_i^{\sigma}(\vec{r}) = \varepsilon_i^{\sigma} \psi_i^{\sigma}(\vec{r})$$

$$V_{eff}^{\sigma}(\vec{r}) = V_{ext}(\vec{r}) + V_{Hartree}[n] + V_{xc}^{\sigma}[n^{\uparrow}, n^{\downarrow}]$$

Goal: solve the equation, that is, **find**

- the **eigenvectors**
- the **eigenvalues**

Solution: expand the eigenvectors in terms of functions of known properties (**basis**)

$$\psi_i(\vec{r}) = \sum_{\alpha} c_{i\alpha} \underbrace{f_{\alpha}(\vec{r})}_{\text{basis functions}}$$

Different methods propose different basis functions

Each method has its own **advantages**:


- most appropriate for a range of problems
- provide insightful information in its realm of application

Each method has its own **pitfalls**:


- importance to understand the method, the pros and the cons.
- what can be computed and what can not be computed

Three main families of methods depending on the basis sets


Atomic sphere methods




Plane wave and grids



Localized basis sets




Order-N methods: The computational load scales linearly with the system size



G. Galli and M. Parrinello, Phys. Rev Lett. 69, 3547 (1992)

Locality is the key point to achieve linear scaling

Large system




x2

"Divide and Conquer"

W. Yang, Phys. Rev. Lett. 66, 1438 (1992)

Efficient basis set for linear scaling calculations: localized, few and confined

Locality \Rightarrow Basis set of **localized functions**




Regarding **efficiency**, the important aspects are:


- **NUMBER** of basis functions per atom
- **RANGE** of localization of these functions

Three main families of methods depending on the basis sets


Atomic sphere methods



Plane wave and grids



Localized basis sets



Atomic orbitals: advantages and pitfalls

$$\phi_{Ilmn}(\vec{r}) = R_{Il}(|\vec{r}|) Y_{lm}(\hat{r})$$

ADVANTAGES

- Very efficient (number of basis functions needed is usually very small).
- Large reduction of CPU time and memory
- Straightforward physical interpretation (population analysis, projected density of states,...)
- They can achieve very high accuracies...

DISADVANTAGES

- ...Lack of systematic for convergence (not unique way of enlarge the basis set)
- Human and computational effort searching for a good basis set before facing a realistic project.
- Depend on the atomic position (Pulay terms).

Atomic orbitals: a radial function times an spherical harmonic


$$\phi_{Iln}(\vec{r}) = R_{Iln}(|\vec{r}_I|) Y_{lm}(\hat{r}_I)$$

Index of an atom \nearrow

Possibility of multiple orbitals with the same l, m \nearrow

Angular momentum \uparrow

$\hat{r}_I = \frac{\vec{r}_I}{|\vec{r}_I|}$




Numerical atomic orbitals

Numerical solution of the Kohn-Sham Hamiltonian for the **isolated pseudoatom** with the **same approximations** (xc,pseudos) as for the condensed system

$$\left(-\frac{1}{2r} \frac{d^2}{dr^2} r + \frac{l(l+1)}{2r^2} + V_l(r) \right) R_l(r) = \varepsilon_l R_l(r)$$

This equation is solved in a logarithmic grid using the Numerov method



Dense close at the origin where
atomic quantities oscillates wildly


Light far away from the origin where
atomic quantities change smoothly

Atomic orbitals:

Main features that characterize the basis

$$\phi_{Ilmn}(\vec{r}) = R_{Ilm}(|\vec{r}_I|) Y_{lm}(\hat{r}_I)$$

Radial part:
degree of freedom to play with



Spherical harmonics:
well defined (fixed) objects



Size: Number of atomic orbitals per atom

Range: Spatial extension of the orbitals


Shape: of the radial part

Size (number of basis set per atom)

Depending on the required accuracy and
available computational power

Quick exploratory
calculations

Highly converged
calculations



Minimal basis set
(single- ζ ; SZ)

Multiple- ζ

+

Polarization

+

Diffuse orbitals

+ Basis optimization

Converging the basis size: from quick and dirty to highly converged calculations

Single- ζ (minimal or SZ)


One single radial function per angular
momentum shell occupied in the free-atom

Examples of minimal basis-set:

Si atomic configuration: $\underbrace{1s^2 2s^2 2p^6}_{\text{core}} \quad \underbrace{3s^2 3p^2}_{\text{valence}}$


$l = 0$ (s)

$m = 0$




$l = 1$ (p)


$m = -1$



$m = 0$



$m = +1$



4 atomic orbitals per Si atom

(pictures courtesy of Victor Luaña)

Converging the basis size: from quick and dirty to highly converged calculations

Single- ζ (minimal or SZ)

One single radial function per angular
momentum shell occupied in the free-atom


Examples of minimal basis-set:

Fe atomic configuration: $\underbrace{1s^2 2s^2 2p^6 3s^2 3p^6}_{\text{core}}$

$\underbrace{4s^2 3d^6}_{\text{valence}}$


$l = 0$ (s)

$m = 0$




$l = 2$ (d)


$m = -2$




$m = -1$




$m = 0$



$m = +1$




$m = +2$




6 atomic orbitals per Fe atom

(pictures courtesy of Victor Luaña)


The optimal atomic orbitals are environment dependent



$R \rightarrow 0$ (He atom)



$R \rightarrow \infty$ (H atom)



Basis set **generated for isolated atoms...** ...
but **used in molecules or condensed systems**

Add flexibility to the basis to adjust to different configurations

Converging the basis size: from quick and dirty to highly converged calculations

Single- ζ (minimal or SZ)

One single radial function per angular
momentum shell occupied in the free-atom




Improving the quality

Radial flexibilization:

Add more than one radial function
within the same angular
momentum than SZ


Multiple- ζ

Default mechanism to generate multiple- ζ in SIESTA: “Split-valence” method




Starting from the function we want to supplement

Default mechanism to generate multiple- ζ in SIESTA: “Split-valence” method



The second- ζ function reproduces the tail of the of the first- ζ outside a radius r_m


Default mechanism to generate multiple- ζ in SIESTA: “Split-valence” method



And continuous smoothly towards the origin as $r^l (a_l - b_l r^2)$


(two parameters: the second- ζ and its first derivative continuous at r_m)

Default mechanism to generate multiple- ζ in SIESTA: “Split-valence” method



The same Hilbert space can be expanded if we use the **difference**, with the advantage that now the second- ζ vanishes at r_m (more efficient)

Default mechanism to generate multiple- ζ in SIESTA: “Split-valence” method



Finally, the second- ζ is normalized

r_m controlled with **PAO.SplitNorm** (typical value 0.15)

Converging the basis size: from quick and dirty to highly converged calculations

Single- ζ (minimal or SZ)

One single radial function per angular
momentum shell occupied in the free-atom

Improving the quality



Radial flexibilization:

Add more than one radial function
within the same angular
momentum than SZ

Multiple- ζ

Angular flexibilization:

Add shells of different atomic
symmetry (different l)

Polarization


Example of adding angular flexibility to an atom

Polarizing the Si basis set

Si atomic configuration: $\underbrace{1s^2 2s^2 2p^6}_{\text{core}} \underbrace{3s^2 3p^2}_{\text{valence}}$


$l = 0$ (s)

$m = 0$




$l = 1$ (p)


$m = -1$



$m = 0$



$m = +1$



Polarize: add $l = 2$ (d) shell






$m = -2$

$m = -1$

$m = 0$

$m = +1$

$m = +2$





New orbitals directed in different directions with respect the original basis

Two different ways of generate polarization orbitals

Perturbative polarization

Apply a **small electric field** to the orbital we want to polarize




Si 3d
orbitals

Two different ways of generate polarization orbitals

Perturbative polarization


Apply a **small electric field** to the orbital we want to polarize



Atomic polarization

Solve **Schrödinger equation** for **higher angular momentum**

unbound in the free atom \Rightarrow
require short cut offs



Si 3d
orbitals


Improving the quality of the basis \Rightarrow more atomic orbitals per atom

Atom	Valence configuration	SZ	DZ	P
		# orbitals symmetry	# orbitals symmetry	# orbitals symmetry
Si	$3s^2 3p^2$	1 s	2 s	1 d_{xy}
		1 p_x	2 p_x	1 d_{yz}
		1 p_y	2 p_y	1 d_{zx}
		1 p_z	2 p_z	1 $d_{x^2-y^2}$
				1 $d_{3z^2-r^2}$
	Total	4	8	(DZ+P) 13


Atom	Valence configuration			
		# orbitals symmetry	# orbitals symmetry	# orbitals symmetry
Fe	$4s^2 3d^6$	1 s	2 s	1 p_x
		1 d_{xy}	2 d_{xy}	1 p_y
		1 d_{yz}	2 d_{yz}	1 p_z
		1 d_{zx}	2 d_{zx}	
		1 $d_{x^2-y^2}$	2 $d_{x^2-y^2}$	
		1 $d_{3z^2-r^2}$	2 $d_{3z^2-r^2}$	
	Total	6	12	(DZ+P) 15

Convergence as a function of the size of the basis set: Bulk Si

Cohesion curves



PW and NAO convergence



Atomic orbitals show nice convergence with respect the size

Polarization orbitals very important for convergence (more than multiple- ζ)

Double- ζ plus polarization equivalent to a PW basis set of 26 Ry

Convergence as a function of the size of the basis set: Bulk Si

	SZ	DZ	TZ	SZP	DZP	TZP	TZDP	PW	APW	Exp
a (Å)	5.52	5.46	5.45	5.42	5.39	5.39	5.39	5.38	5.41	5.43
B (GPa)	89	96	98	98	97	97	96	96	96	98.8
E _c (eV)	4.72	4.84	4.91	5.23	5.33	5.34	5.34	5.37	5.28	4.63

A DZP basis set introduces the same deviations as the ones due to the DFT or the pseudopotential approaches

SZ = single- ζ

P=Polarized

PW: Converged Plane Waves (50 Ry)

DZ= double- ζ


DP=Doble-
polarized

APW: Augmented Plane Waves

TZ=triple- ζ

Range: the spatial extension of the atomic orbitals

Order(N) methods \Rightarrow locality, that is, a finite range for matrix and overlap matrices



If the two orbitals are sufficiently far away

$$S_{\nu\mu} = \langle \phi_\nu | \phi_\mu \rangle = \int d\vec{r} \phi_\nu^*(\vec{r}) \phi_\mu(\vec{r}) = 0$$

$$H_{\nu\mu} = \langle \phi_\nu | \hat{H} | \phi_\mu \rangle = \int d\vec{r} \phi_\nu^*(\vec{r}) \hat{H} \phi_\mu(\vec{r}) = 0$$

Neglect interactions:

Below a tolerance

Beyond a given scope of neighbours

Difficulty: introduce numerical instabilities for high tolerances.

Strictly localized atomic orbitals:


Vanishes beyond a given cutoff radius

O. Sankey and D. Niklewski, PRB 40, 3979 (89)

Difficulty: accuracy and computational efficiency depend on the range of the basis orbitals

How to define all the r_c in a balance way?

How to control the range of the orbitals in a balanced way: the energy shift



Particle in a confinement potential:

Imposing a finite $\int_{-\infty}^{+\infty} |\phi(x)|^2 dx$
+

Continuous function and first derivative



E is quantized (not all values allowed)

Increasing $E \Rightarrow \phi_\mu$ has a node
and tends to $-\infty$ when $x \rightarrow -\infty$

Complement M III "Quantum Mechanics",
C. Cohen-Tannoudji *et al.*

How to control the range of the orbitals in a balanced way: the energy shift

$$\left(-\frac{1}{2r} \frac{d^2}{dr^2} r + \frac{l(l+1)}{2r^2} + V_l(r) \right) R_l(r) = (\varepsilon_l + \delta\varepsilon_l) R_l(r)$$



Energy increase = Energy shift

PAO.EnergyShift (energy)


Cutoff radius, r_c , = position where each orbital has the node

A **single parameter for all** cutoff radii

The larger the Energy shift, the shorter the r_c 's

Typical values: 100-200 meV

Convergence with the range



Bulk Si

equal s, p
orbitals radii

J. Soler *et al.*, J. Phys: Condens. Matter, 14, 2745 (2002)

More efficient



More accurate

The range and shape might be also controlled by an extra charge and/or by a confinement potential

Extra charge δQ

Orbitals in **anions** tend to be more **delocalized**

Orbitals in **cations** tend to be more **localized**

(For instance, this parameter might be important in some oxides)

Confinement potentials


Solve the Schrödinger equation for the isolated atom inside an confinement potential

Different proposals for the confinement potentials: Hard confinement

Fireball

O. F. Sankey and D. J. Niklewski, Phys. Rev. B 40, 3979 (89)


The default in SIESTA



Determined by the energy shift

Advantages: empirically, it works very nice

Pitfall: produces orbitals with first derivative discontinuous at r_c
problem when combined with numerical grids.




Different proposals for the confinement potentials: Soft-confinement potential

Available in SIESTA

$$V(r) = V_0 \frac{e^{-\frac{r_c - r_i}{r - r_i}}}{r_c - r}$$

J. Junquera *et al.*, Phys. Rev. B 64, 235111 (2001)



Advantages:

- orbital continuous with all the derivatives continuous
- diverges at r_c (orbital exactly vanishes there)
- zero at the core region

Pitfall: two new parameters to play with, more exploratory calculations

How to introduce the basis set in SIESTA

Effort on defining a systematic with minimum parameters

If **nothing** is specified: **default**

		Default value
Basis size:	PAO.BasisSize	DZP
Range of first-zeta:	PAO.EnergyShift	0.02 Ry
Second-zeta:	PAO.BasisType	Split
Range of second-zeta:	PAO.SplitNorm	0.15
Confinement:	Hard well	

Good basis set in terms of accuracy versus efficiency

More global control on the basis with a few input variables: size and range

Size:

Basis size:

PAO.BasisSize

SZ

DZ

SZP

DZP

Range:

Range of first-zeta: PAO.EnergyShift 0.02 Ry

Range of second-zeta: PAO.SplitNorm 0.15

The larger both values, the more confined the basis functions


More specific control on the basis: the PAO.Basis block



More specific control on the basis: the PAO.Basis block

Some variable might be computed automatically

```
%block PAO.Basis                                # Define Basis set
H          1    +0.25                             # Species label, number of l-shells, charge
n=1        0    2                                # n, l, Nzeta
0.000      0.000                                # rc (first-zeta), rm (second-zeta)
1.000      1.000                                # scaling factors
%endblock PAO.Basis
```



These variables calculated from
PAO.EnergyShift and PAO.SplitNorm values

More specific control on the basis: the PAO.Basis block

Adding polarization orbitals: perturbative polarization

```
%block PA0.Basis
H      1    +0.25
n=1    0    2 P    1
      5.000      3.000
      1.000      1.000
%endblock PA0.Basis
```

**More specific control on the basis:
the PAO.Basis block**

Adding polarization orbitals: atomic polarization

**More specific control on the basis:
the PAO.Basis block**

Adding polarization orbitals: atomic polarization

[illegible]

More specific control on the basis: the PAO.Basis block

Soft-confinement potential

```
%block PAO.Basis                                # Define Basis set
H          1    +0.25                             # Species label, number of l-shells, charge
n=1        0    2 E 150.00 4.5                   # n, l, Nzeta, flag soft-conf, prefactor, inner rad
          5.000    3.000                         # rc (first-zeta), rm (second-zeta)
          1.000    1.000                         # scaling factors
%endblock PAO.Basis
```

$$V(r) = V_0 \frac{e^{-\frac{r_c - r_i}{r - r_i}}}{r_c - r}$$

V_0 in Ry

r_i in bohrs

Recap

Numerical Atomic Orbitals

A very efficient basis set

Especially suitable for Order-N methods

Smooth transition from quick exploratory calculations to highly converged

Lack of systematic convergence

Simple handles for tuning the basis sets

Generate multiple- ζ : Split Valence

Generate polarization orbitals: Perturbative polarization

Control the range of the orbitals in a balanced way: Energy Shift

Confine the orbitals: Soft-confinement potential

A DZP basis set, the same deviations as DFT functional or Pseudo

Supplementary information

Spherical Bessel functions $j_l(kr)$, solutions of a free particle confined in a box



$$V = \infty, r \geq a$$


Schrödinger equation for a particle inside the box

$$-\frac{\hbar^2}{2m} \nabla^2 \psi(r, \theta, \phi) = E \psi(r, \theta, \phi)$$

After separation of variables, the radial equation reads

$$\psi(r, \theta, \phi) = R(r) \Theta(\theta) \Phi(\phi)$$

$$V = 0, r < a$$



$$\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + \left[k^2 - \frac{l(l+1)}{r^2} \right] R = 0$$

$$k^2 = \frac{2mE}{\hbar^2}$$

$l \in \mathbf{Z}$, separation
variable constant

Solution of the radial equation

$$R(r) = \begin{cases} A j_l(kr) + B n_l(kr), & r < a \\ 0, & r \geq a \end{cases}$$

Boundary conditions: k must satisfy $j_l(ka) = 0$

Spherical von Neumann
function, not finite at the origin